

VISUALIZING ELEMENTS OF SHA[3] IN GENUS 2 JACOBIANS

NILS BRUIN AND SANDER R. DAHMEN

ABSTRACT. Mazur proved that any element ξ of order three in the Shafarevich-Tate group of an elliptic curve E over a number field k can be made visible in an abelian surface A in the sense that ξ lies in the kernel of the natural homomorphism between the cohomology groups $H^1(\text{Gal}(\bar{k}/k), E) \rightarrow H^1(\text{Gal}(\bar{k}/k), A)$. However, the abelian surface in Mazur's construction is almost never a jacobian of a genus 2 curve. In this paper we show that any element of order three in the Shafarevich-Tate group of an elliptic curve over a number field can be visualized in the jacobians of a genus 2 curve. Moreover, we describe how to get explicit models of the genus 2 curves involved.

1. INTRODUCTION

Let E be an elliptic curve over a field k with separable closure $\bar{k}$. We write $H^1(k, E[3]) := H^1(\text{Gal}(\bar{k}/k), E[3](\bar{k}))$ for the first galois cohomology group taking values in the 3-torsion of E (the notation $H^i(k, A)$ is used similarly for other group schemes A/k later in this paper). We are primarily concerned with the question which $\delta \in H^1(k, E[3])$ are *visible* in the jacobian of a genus 2 curve. Mazur defines *visibility* in the following way. Let $0 \rightarrow E \rightarrow A \rightarrow B \rightarrow 0$ be a short exact sequence of abelian varieties over k . By taking galois cohomology, we obtain the exact sequence

$$(1.1) \quad A(k) \longrightarrow B(k) \longrightarrow H^1(k, E) \xrightarrow{\phi} H^1(k, A) .$$

Elements of the kernel of ϕ are said to be *visible* in A . Mazur chose this term because a model of the principal homogeneous space corresponding to an element $\xi \in H^1(k, E)$ that is visible in A can be obtained as a fiber of A over a point in $B(k)$ (this can readily be seen from (1.1)). By extension, we say that $\delta \in H^1(k, E[n])$ is visible in A if the image of δ under the natural homomorphism $H^1(k, E[n]) \rightarrow H^1(k, E)$ is visible in A .

Let us restrict to the case that k is a number field for the rest of this section. Inspired by some surprising experimental data [2], Mazur [3] proved, that for any element ξ in the Shafarevich-Tate group $\text{III}(E/k)$ of order three, there exists an abelian variety A over k such that ξ is visible in A . The abelian variety that Mazur constructs is almost never principally polarizable over $\bar{k}$ and hence is almost never a jacobian of a genus 2 curve. In the present paper, we show that any element from $\text{III}(E/k)[3]$ is in fact visible in the jacobian of a genus 2 curve. Moreover, we describe how to get an explicit model of such a genus 2 curve.

Date: January 28, 2010.

2000 Mathematics Subject Classification. Primary 11G30; Secondary 14H40.

Key words and phrases. Elliptic curves, Visibility, Shafarevich-Tate groups, Weil pairing.

Research of both authors supported by NSERC.

2. TORSORS AND THETA GROUPS

Throughout this section let $n > 1$ be an integer, let k be a perfect field of characteristic not dividing n and let E denote an elliptic curve over k . In [1], many equivalent interpretations are given for the group $H^1(k, E[n])$. For our purposes, we need two classes of objects. The first is most closely related with descent in general and our question in particular. We consider E -torsors under $E[n](\bar{k})$ and, following [1], call them n -coverings.

Definition 2.1. An n -covering $\pi : C \rightarrow E$ of an elliptic curve E is an unramified covering over k that is galois and irreducible over $\bar{k}$, with $\text{Aut}_{\bar{k}}(C/E) \simeq E[n](\bar{k})$. Two n -coverings $\pi_1 : C_1 \rightarrow E$, $\pi_2 : C_2 \rightarrow E$ are called isomorphic if there exists a k -morphism $\phi : C_1 \rightarrow C_2$ such that $\pi_1 = \pi_2 \circ \phi$.

Over $\bar{k}$, all n -coverings are isomorphic to the trivial n -covering, the multiplication-by- n map $[n] : E \rightarrow E$.

Proposition 2.2 ([1, Proposition 1.14]). *The k -isomorphism classes of n -coverings of E are classified by $H^1(k, E[n])$.*

For $\delta \in H^1(k, E[n])$ we denote by C_δ the curve in the covering $C_\delta \rightarrow E$ corresponding to δ . We remark that $\delta \in H^1(k, E[n])$ has trivial image in $H^1(k, E)$ if and only if C_δ has a k -rational point.

We write O for the identity on E . The complete linear system $|n \cdot O|$ determines a morphism $E \rightarrow \mathbb{P}^{n-1}$, where the translation action of $E[n]$ extends to a linear action on $\mathbb{P}^{n-1}$. This gives a projective representation $E[n] \rightarrow \text{PGL}_n$. The lift of this representation to GL_n gives rise to a group Θ_E , which fits in the following diagram.

$$(2.1) \quad \begin{array}{ccccccc} 1 & \longrightarrow & \mathbb{G}_m & \xrightarrow{\alpha_E} & \Theta_E & \xrightarrow{\beta_E} & E[n] \longrightarrow 1 \\ & & \parallel & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \mathbb{G}_m & \longrightarrow & \text{GL}_n & \longrightarrow & \text{PGL}_n \longrightarrow 1 \end{array}$$

The group $E[n](\bar{k})$ carries additional structure. It also has the Weil pairing e_E , which is a non-degenerate alternating galois covariant pairing taking values in the n -th roots of unity

$$e_E : E[n](\bar{k}) \times E[n](\bar{k}) \rightarrow \mu_n(\bar{k}).$$

The commutator of Θ_E corresponds to the Weil pairing, meaning that for $x, y \in \Theta_E$ we have

$$xyx^{-1}y^{-1} = \alpha_E(e_E(\beta_E(x), \beta_E(y))).$$

Definition 2.3. A theta group for $E[n]$ is a central extension of group schemes

$$1 \rightarrow \mathbb{G}_m \xrightarrow{\alpha} \Theta \xrightarrow{\beta} E[n] \rightarrow 1$$

such that the Weil-pairing on $E[n]$ corresponds to the commutator, i.e. for $x, y \in \Theta$ we have

$$xyx^{-1}y^{-1} = \alpha(e_E(\beta(x), \beta(y))).$$

Two theta groups

$$1 \rightarrow \mathbb{G}_m \rightarrow \Theta_i \rightarrow E[n] \rightarrow 1, \quad i = 1, 2$$

are called isomorphic if there exists a group scheme isomorphism $\phi : \Theta_1 \rightarrow \Theta_2$ over k making the following diagram commutative.

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathbb{G}_m & \longrightarrow & \Theta_1 & \longrightarrow & E[n] \longrightarrow 1 \\ & & \parallel & & \downarrow \phi & & \parallel \\ 1 & \longrightarrow & \mathbb{G}_m & \longrightarrow & \Theta_2 & \longrightarrow & E[n] \longrightarrow 1 \end{array}$$

Over $\bar{k}$, all theta-groups are isomorphic to Θ_E as central extensions; see [1, Lemma 1.30].

Proposition 2.4 ([1, Proposition 1.31]). *Let $E[n]$ be the n -torsion subscheme of an elliptic curve E over a field k , equipped with its Weil pairing. The isomorphism classes of theta-groups for $E[n]$ over k are classified by $H^1(k, E[n])$.*

The theta group associated to $\delta \in H^1(k, E[n])$ may allow for a matrix representation $\Theta \rightarrow \mathrm{GL}_n$ that fits in a diagram like (2.1). This is measured by the *obstruction map* Ob introduced in [4] and [1]. This map can be obtained by taking non-abelian galois cohomology of the defining sequence of Θ_E :

$$\cdots \longrightarrow H^1(k, \Theta_E) \longrightarrow H^1(k, E[n]) \xrightarrow{\mathrm{Ob}} H^2(k, \mathbb{G}_m) = \mathrm{Br}(k) \longrightarrow \cdots$$

Note that, except in some trivial cases, Ob is *not* a group homomorphism. The map Ob also has an interpretation in terms of n -coverings. Let $C \rightarrow E$ be an n -covering associated to $\delta \in H^1(k, E[n])$. We have that $\mathrm{Ob}(\delta) = 0$ if and only if C admits a model $C \rightarrow \mathbb{P}^{n-1}$ with $\mathrm{Aut}_{\bar{k}}(C/E) = E[3](\bar{k})$ acting linearly, in which case C is $\bar{k}$ -isomorphic to E as a curve and the covering $C \rightarrow E$ is simply a translation composed with multiplication-by- n .

Remark 2.5. Note that if k is a number field, then any element in $\mathrm{Br}(k)$ that restricts to the trivial element in $\mathrm{Br}(k_v)$ in all completions k_v of k , is trivial itself. It follows that Ob is trivial on the n -Selmer group $S^{(n)}(E/k)$.

3. VISIBILITY IN SURFACES

Let E_1 be an elliptic curve over a perfect field k of characteristic distinct from 3. In what follows, we will consider $\delta \in H^1(k, E_1[3])$ with $\mathrm{Ob}(\delta) = 0$. A possible way of constructing an abelian surface A such that δ is visible in A starts by taking a suitable elliptic curve E_2/k together with a k -group scheme isomorphism $\lambda : E_1[3] \rightarrow E_2[3]$. Let $\Delta \subset E_1 \times E_2$ be the graph of $-\lambda$ so that

$$\Delta(\bar{k}) = \{(P, -\lambda(P)) : P \in E_1[3](\bar{k})\}.$$

Let $A := (E_1 \times E_2)/\Delta$ and write $\phi : E_1 \times E_2 \rightarrow A$ for the corresponding isogeny. Since $\Delta \subset E_1[3] \times E_2[3]$, we have another isogeny $\phi' : A \rightarrow E_1 \times E_2$ such that $\phi' \circ \phi = 3$. We write p^* for the composition $E_1 \rightarrow (E_1 \times E_2) \xrightarrow{\phi} A$ and p_* for the composition $A \xrightarrow{\phi'} (E_1 \times E_2) \rightarrow E_1$ and q^*, q_* for the corresponding morphisms concerning E_2 . It is straightforward to verify that p^*, q^* are embeddings, that $\phi = p^* + q^*$ (where the projections are understood and we note that the $+$ sign here corresponds to the $-$ sign in the definition of Δ) and that $\phi' = p_* \times q_*$.

We combine the galois cohomology of the short exact sequences

$$\begin{aligned} 0 \rightarrow E_1 \xrightarrow{p^*} A \xrightarrow{q^*} E_2 \rightarrow 0, \\ 0 \rightarrow E_2 \xrightarrow{q^*} A \xrightarrow{p^*} E_1 \rightarrow 0, \text{ and} \\ 0 \rightarrow E_i[3] \rightarrow E_i \xrightarrow{3} E_i \rightarrow 0 \text{ for } i = 1, 2 \end{aligned}$$

to obtain the big (symmetric) commutative diagram with exact rows and columns

$$\begin{array}{ccccccc} & & & & E_2(k) & \xrightarrow{q^*} & A(k) \\ & & & & \downarrow 3 & & \downarrow q_* \\ & & & & E_2(k) & \xlongequal{\quad} & E_2(k) \\ & & & & \downarrow \alpha & & \downarrow \\ E_1(k) & \xrightarrow{3} & E_1(k) & \longrightarrow & H^1(k, \Delta) & \longrightarrow & H^1(k, E_1) \\ \downarrow p^* & & \parallel & & \downarrow & & \downarrow \\ A(k) & \xrightarrow{p_*} & E_1(k) & \longrightarrow & H^1(k, E_2) & \longrightarrow & H^1(k, A) \end{array}$$

where we note that $H^1(k, \Delta) \simeq H^1(k, E_1[3]) \simeq H^1(k, E_2[3])$. We see that δ is visible in A precisely if $\delta \in H^1(k, E_1[3]) = H^1(k, \Delta)$ lies in the image of α , i.e., if the curve $C_{\lambda(\delta)}$ corresponding to $\lambda(\delta) \in H^1(k, E_2[3])$ has a rational point. We summarize these observations, which are due to Mazur.

Lemma 3.1. *Let E_1 be an elliptic curve over a perfect field k of characteristic distinct from 3 and let $\delta \in H^1(k, E[3])$ with $\text{Ob}(\delta) = 0$. Suppose that there exists an elliptic curve E_2/k and a k -group scheme isomorphism $\lambda : E_1[3] \rightarrow E_2[3]$ such that the curve $C_{\lambda(\delta)}$ corresponding to $\lambda(\delta)$ has a k -rational point. Then δ is visible in the abelian surface $(E_1 \times E_2)/\Delta$ where Δ denotes the graph of $-\lambda$.*

Mazur also observed, in the case of a number field k , that if $\delta \in S^{(3)}(E/k)$, then C_δ admits a plane cubic model. Furthermore, there is a pencil of cubics through the 9 flexes of C_δ , and each non-singular member corresponds to a 3-covering $C_t \rightarrow E_t$, where $E_t[3] \simeq E[3]$ and $C_t \rightarrow E_t$ represents δ . It is therefore easy to find a t such that C_t has a rational point; simply pick a rational point and solve for t . To refine the construction, one can ask

Question 3.2. Can one make $\delta \in H^1(k, E[3])$ visible in the jacobian of a genus 2 curve?

Note that $E_1 \times E_2$ is principally polarized via the product polarization. This gives rise to a Weil pairing on $(E_1 \times E_2)[3]$, corresponding to the product pairing. If A is a jacobian, then A must be principally polarized over $\bar{k}$. One way this could happen is if the isogeny $p^* + q^* : E_1 \times E_2 \rightarrow A$ gives rise to a principal polarization. This would be the case if the kernel Δ is a maximal isotropic subgroup of $E_1[3] \times E_2[3]$ with respect to the product pairing. That means that $\lambda : E_1[3] \rightarrow E_2[3]$ must be an *anti*-isometry, i.e. for all $P, Q \in E_1[3]$ we must have

$$e_{E_2}(\lambda(P), \lambda(Q)) = e_{E_1}(P, Q)^{-1}.$$

Note that the original cubic C is a member of the pencil that Mazur constructs, so in his construction λ is actually an *isometry*, i.e. it preserves the Weil-pairing. Below we consider a pencil of cubics that leads to an anti-isometry λ .

4. ANTI-ISOMETRIC PENCILS

Let k be a perfect field of characteristic distinct from 2, 3. Following [5], we associate to a ternary cubic form $F \in k[x, y, z]$ three more ternary cubic forms. Namely, the Hessian of F

$$H(F) := -\frac{1}{2} \begin{vmatrix} \frac{\partial^2 F}{\partial x \partial x} & \frac{\partial^2 F}{\partial x \partial y} & \frac{\partial^2 F}{\partial x \partial z} \\ \frac{\partial^2 F}{\partial y \partial x} & \frac{\partial^2 F}{\partial y \partial y} & \frac{\partial^2 F}{\partial y \partial z} \\ \frac{\partial^2 F}{\partial z \partial x} & \frac{\partial^2 F}{\partial z \partial y} & \frac{\partial^2 F}{\partial z \partial z} \end{vmatrix},$$

the Caylean of F

$$P(F) := -\frac{1}{xyz} \begin{vmatrix} \frac{\partial F}{\partial x}(0, z, -y) & \frac{\partial F}{\partial y}(0, z, -y) & \frac{\partial F}{\partial z}(0, z, -y) \\ \frac{\partial F}{\partial x}(-z, 0, x) & \frac{\partial F}{\partial y}(-z, 0, x) & \frac{\partial F}{\partial z}(-z, 0, x) \\ \frac{\partial F}{\partial x}(y, -x, 0) & \frac{\partial F}{\partial y}(y, -x, 0) & \frac{\partial F}{\partial z}(y, -x, 0) \end{vmatrix}$$

and a ternary cubic form denoted $Q(F)$, for which we refer to [5, Section 11.2]. For most cases one can take $Q(F)$ to be $H(P(F))$ or $P(H(F))$, but there are some exceptional cases where $P(F), Q(F)$ span an appropriate pencil and $P(F), H(P(F))$ do not. The left action of GL_3 on k^3 induces a right action of GL_3 on ternary cubic forms (or, more generally, on $k[x, y, z]$). For a ternary cubic form F and an $M \in \mathrm{GL}_3$ we denote this action simply by $F \circ M$. The significance of the three associated ternary cubic forms lies in the fact that $H(F)$ depends covariantly on F (of weight 2) and $P(F)$ and $Q(F)$ depend contravariantly on F (of weights 4 and 6 respectively). This means that for every ternary cubic form F and every $M \in \mathrm{GL}_3$ we have, with $d := \det M$ that

$$\begin{aligned} H(F \circ M) &= d^2 H(F) \circ M \\ P(F \circ M) &= d^4 P(F) \circ M^{-T} \\ Q(F \circ M) &= d^6 Q(F) \circ M^{-T}, \end{aligned}$$

where M^{-T} denotes the inverse transpose of M .

Now consider a smooth cubic curve C in $\mathbb{P}^2$ given by the zero locus of a ternary cubic form F . Then C has exactly 9 different flex points Φ , which all lie on the (not necessarily smooth) curve given by $H(F) = 0$. The smoothness of C guarantees that F and $H(F)$ will be linearly independent over k . Hence Φ can be described as the intersection $F = H(F) = 0$. We call Φ the *flex scheme* of C . At least one of $P(F)$ and $Q(F)$ turns out to be nonsingular (still assuming that C is nonsingular) and the intersection $P(F) = Q(F) = 0$ equals the flex points Φ^* of the nonsingular cubics among $P(F)$ and $Q(F)$ (if, say, $P(F)$ is nonsingular, then Φ^* can of course also be written as $P(F) = H(P(F)) = 0$).

We can consider the pencil of cubics through Φ , explicitly given by

$$C_{(s:t)} : sF(x, y, z) + tH(F)(x, y, z) = 0.$$

Classical invariant theory tells us the following. This pencil has exactly 4 singular members and all other members have flex scheme equal to Φ . Conversely, any nonsingular cubic with flex scheme Φ occurs in this pencil. Furthermore, both $P(sF + tH(F))$ and $Q(sF + tH(F))$ are linear combinations of $P(F)$ and $Q(F)$.

This shows that the flex scheme Φ^* is independent of the choice of C through Φ and only depends on Φ . We call Φ^* the dual flex scheme of Φ and we will justify this name below.

Remark 4.1. In the discussion above it was convenient to consider just one projective plane $\mathbb{P}^2$. A more canonical way would be to consider a projective plane $\mathbb{P}^2$ with coordinates x, y, z and the dual projective plane, denoted $(\mathbb{P}^2)^*$, whose coordinates u, v, w are related to those of $\mathbb{P}^2$ by $ux + vy + wz = 0$. Now let C be a smooth cubic curve in $\mathbb{P}^2$ given by the zero locus of the ternary cubic form $F(x, y, z)$ with flex scheme Φ . The 9 tangent lines through Φ determine 9 points in $(\mathbb{P}^2)^*$. Generically, these 9 points in $(\mathbb{P}^2)^*$ will not be the flex points of a smooth cubic curve, hence generically there will be a unique cubic curve going through these points. This curve in $(\mathbb{P}^2)^*$ is exactly given by the zero locus of the Caylean, i.e. $P(F)(u, v, w) = 0$; see also [6, pp.151,190–191]. Moreover, if the characteristic of k is zero, then it turns out that this cubic curve is nonsingular if and only if the j -invariant of C is nonzero.

As a simple, but important example we take $F := x^3 + y^3 + z^3$. Then we compute

$$H(F) = -108xyz, \quad P(F) = -54xyz, \quad Q(F) = 324(x^3 + y^3 + z^3).$$

Now define Φ_0 to be the flex scheme of $F = 0$, i.e.

$$(4.1) \quad \Phi_0 := \{[x : y : z] \in \mathbb{P}^2 : x^3 + y^3 + z^3 = xyz = 0\}.$$

Then we see that the flex scheme given by $P(F) = Q(F) = 0$ (which is the flex scheme of $Q(F) = 0$) equals Φ_0 , i.e.

$$\Phi_0^* = \Phi_0.$$

Geometrically all flex schemes are linear transformations of each other. In particular, for any flex scheme Φ there exists an $M \in \mathrm{GL}_3(\bar{k})$ such that $\Phi = M\Phi_0$. The contravariance of P and Q implies that the assignment $\Phi \mapsto \Phi^*$ has the contravariance property that for any flex scheme Φ and $M \in \mathrm{GL}_3$

$$(4.2) \quad (M\Phi)^* = M^{-T}\Phi^*.$$

We also note that this implies that the assignment $\Phi \mapsto \Phi^{**} := (\Phi^*)^*$ is covariant in the sense that for any flex scheme Φ and $M \in \mathrm{GL}_3$ we have $(M\Phi)^{**} = M\Phi^{**}$. Writing $\Phi = M\Phi_0$ and using $(\Phi_0)^{**} = \Phi_0^* = \Phi_0$ we now get

$$\Phi^{**} = (M\Phi_0)^{**} = M\Phi_0^{**} = M\Phi_0 = \Phi.$$

This justifies calling Φ^* the *dual* flex scheme of Φ .

To any flex scheme Φ we associate a group $\Theta(\Phi) \subset \mathrm{GL}_3$ as follows. Choose a nonsingular cubic curve C through Φ and let E be its jacobian. After identifying E and C as curves over $\bar{k}$, we get an action of $E[3]$ on C , which extends to a linear action on $\mathbb{P}^2$. This determines an embedding $\chi : E[3] \rightarrow \mathrm{PGL}_3$. Obviously, the image $\chi(E[3])$ only depends on Φ . We define $\Theta(\Phi)$ to be the inverse image of $\chi(E[3])$ in GL_3 . Actually $\Theta(\Phi)$ can be defined just in terms of Φ , without choosing C , since it turns out that $\chi(E[3])$ consists exactly of the linear transformations that preserve Φ . (One way of quickly finding these linear transformations explicitly is by using the fact that, for any two distinct points of Φ , the line through these two points intersects Φ in a unique third point.) The construction gives rise to the theta group

$$1 \rightarrow \mathbb{G}_m \rightarrow \Theta(\Phi) \rightarrow E[3] \rightarrow 1.$$

Note that the isomorphism class of this theta group may still depend on the choice of identification of C with E . This corresponds to the choice of an isomorphism between $\Theta(\Phi)/\mathbb{G}_m$ and $E[3]$. If Φ is defined over k , then $E[3]$ and $\Theta(\Phi)$ are also defined over k and the element in $H^1(k, E[3])$ corresponding to this theta group is the same as the element corresponding to the 3-covering $C \rightarrow C/E[3] \simeq E$ for any nonsingular cubic curve C through Φ . The construction also shows that for any $M \in \mathrm{GL}_3$ we have

$$(4.3) \quad \Theta(M\Phi) = M\Theta(\Phi)M^{-1}.$$

Proposition 4.2. *Let $\Phi_1 \subset \mathbb{P}^2$ be a flex scheme and let $\Phi_2 := \Phi_1^*$ be the dual flex scheme. For $i = 1, 2$ let C_i be a smooth plane cubic with flex scheme Φ_i , denote its jacobian by E_i and consider an induced theta group*

$$(4.4) \quad 1 \longrightarrow \mathbb{G}_m \xrightarrow{\alpha_i} \Theta(\Phi_i) \xrightarrow{\beta_i} E_i[3] \longrightarrow 1.$$

Then the outer automorphism $(-T) : \mathrm{GL}_3 \rightarrow \mathrm{GL}_3$ given by $M \mapsto M^{-T}$, yields an isomorphism $\Theta(\Phi_1) \rightarrow \Theta(\Phi_2)$. There exists an anti-isometry $\lambda : E_1[3] \rightarrow E_2[3]$ making the following diagram commutative.

$$(4.5) \quad \begin{array}{ccccccc} 1 & \longrightarrow & \mathbb{G}_m & \xrightarrow{\alpha_1} & \Theta(\Phi_1) & \xrightarrow{\beta_1} & E_1[3] \longrightarrow 1 \\ & & \downarrow x \mapsto x^{-1} & & \downarrow (-T) & & \downarrow \lambda \\ 1 & \longrightarrow & \mathbb{G}_m & \xrightarrow{\alpha_2} & \Theta(\Phi_2) & \xrightarrow{\beta_2} & E_2[3] \longrightarrow 1 \end{array}$$

In particular, let $\delta_i \in H^1(k, E_i[3])$ correspond to the theta group (4.4). Then under the isomorphism $H^1(k, E_1[3]) \simeq H^1(k, E_2[3])$ induced by λ , the cocycle δ_1 maps to δ_2 .

Proof. Once the isomorphism $\Theta(\Phi_1) \rightarrow \Theta(\Phi_2)$ given by $M \mapsto M^{-T}$ is established, the existence of an isomorphism $\lambda : E_1[3] \rightarrow E_2[3]$ making the diagram (4.5) commutative, follows immediately. That λ must be an anti-isometry can readily be seen as follows. Let $P, Q \in E_1[3]$ and choose $x, y \in \Theta(\Phi_1)$ such that $P = \beta_1(x)$ and $Q = \beta_1(y)$. Then

$$\begin{aligned} \alpha_2(e_{E_2}(\lambda(P), \lambda(Q))) &= \alpha_2(e_{E_2}(\beta_2(x^{-T}), \beta_2(y^{-T}))) \\ &= x^{-T} y^{-T} x^T y^T \\ &= (xyx^{-1}y^{-1})^{-T} \\ &= \alpha_1(e_{E_1}(\beta_1(x), \beta_1(y)))^{-T} \\ &= \alpha_1(e_{E_1}(P, Q))^{-1}. \end{aligned}$$

The last statement of the proposition is also immediate, so we are left with establishing $(-T) : \Theta(\Phi_1) \xrightarrow{\sim} \Theta(\Phi_2)$. It suffices to show that for a flex scheme $\Phi \subset \mathbb{P}^2$ we have $\Theta(\Phi)^{-T} = \Theta(\Phi^*)$. Write $\Phi = M\Phi_0$ for some $M \in \mathrm{GL}_3$ with Φ_0 given by (4.1). Then a straightforward calculation shows that $\Theta(\Phi_0)^{-T} = \Theta(\Phi_0)$. We also know that $\Phi_0^* = \Phi_0$, so we get $\Theta(\Phi_0)^{-T} = \Theta(\Phi_0^*)$. Together with (4.2) and (4.3)

we finally obtain,

$$\begin{aligned}
\Theta(\Phi)^{-T} &= \Theta(M\Phi_0)^{-T} \\
&= M^{-T}\Theta(\Phi_0)^{-T}M^T \\
&= M^{-T}\Theta(\Phi_0^*)(M^{-T})^{-1} \\
&= \Theta(M^{-T}\Phi_0^*) \\
&= \Theta((M\Phi_0)^*) \\
&= \Theta(\Phi^*).
\end{aligned}$$

□

Remark 4.3. The construction above of the dual flex scheme Φ^* of a flex scheme Φ involved choosing a smooth cubic going through Φ . Without using theta groups, it was not obvious from this construction that the degree 9 étale algebra $k(\Phi)$ is isomorphic to $k(\Phi^*)$. However, there exists a nice explicit geometric construction of the dual flex scheme that remedies these shortcomings of the earlier construction. Given a flex scheme Φ , we proceed as follows. We label its 9 points over $\bar{k}$ with $P_1, \dots, P_9$. There are 4 sets of 3 lines, (corresponding to the 4 singular member of the pencil of cubics through ϕ) containing these points. We label the line that contains P_i, P_j, P_k with $l_{\{i,j,k\}}$. One can label the points such that the subscripts are

$$\begin{array}{cccc}
\{1, 2, 3\} & \{1, 4, 7\} & \{1, 5, 9\} & \{1, 6, 8\} \\
\{4, 5, 6\} & \{2, 5, 8\} & \{2, 6, 7\} & \{2, 4, 9\} \\
\{7, 8, 9\} & \{3, 6, 9\} & \{3, 4, 8\} & \{3, 5, 7\}
\end{array}$$

Naturally, two different lines $l_{\{i_1, j_1, k_1\}}, l_{\{i_2, j_2, k_2\}}$ meet in a unique point. If for example $i_1 = i_2$, then the intersection point is P_{i_1} . If the two sets $\{i_1, j_1, k_1\}, \{i_2, j_2, k_2\}$ are disjoint, then the two lines meet in a point outside Φ . We name this point $L_{\{i_3, j_3, k_3\}}$, where $\{i_1, j_1, k_1, i_2, j_2, k_2, i_3, j_3, k_3\} = \{1, \dots, 9\}$. As it turns out, the four points that have i in their label all lie on a line p_i . It is also straightforward to check that the p_i together with the $L_{\{i,j,k\}}$ form a configuration in $(\mathbb{P}^2)^*$ that is completely dual to the P_i with the $l_{\{i,j,k\}}$. The p_i form the $\bar{k}$ points of a flex scheme in $(\mathbb{P}^2)^*$, which is justifiably a flex scheme Φ^* dual to Φ , and its construction immediately implies the contravariance property $(M\Phi)^* = M^{-T}\Phi^*$.

We can easily verify that the two constructions of Φ^* coincide for one flex scheme, for instance Φ_0 . The general result then follows because any flex scheme can be expressed as $M\Phi_0$ for some $M \in \text{GL}_3(\bar{k})$.

Since the action of $\text{Gal}(\bar{k}/k)$ on $\{P_1, \dots, P_9\}$ must act via collinearity-preserving permutations, we see that if $\sigma(P_i) = P_{\sigma(i)}$ then $\sigma(p_i) = p_{\sigma(i)}$. Hence, we see that the $\bar{k}$ -points of Φ and its dual have the same Galois action and hence $k(\Phi)$ is isomorphic as a k -algebra to $k(\Phi^*)$.

5. RECOVERING THE GENUS 2 CURVE

Let k be a field and let E_1, E_2 be two elliptic curves over k with an anti-isometry $\lambda : E_1[3] \rightarrow E_2[3]$ and denote by Δ the graph of $-\lambda$ as before. Recall that $E_1 \times E_2$ is principally polarized via the product polarization and that the induced polarization on $A := (E_1 \times E_2)/\Delta$ is also principal in this case. It is a classical fact that if A is not geometrically isomorphic to a product of elliptic curves, then A (together with its principal polarization) is isomorphic to the jacobian of a genus 2 curve C . Let

us assume from now on that E_1 and E_2 are non-isogenous. In [7] it is shown that in this case A is always isomorphic over k to the jacobian of a genus 2 curve C/k . This is enough to get our main theoretical result.

Theorem 5.1. *Let E be an elliptic curve over a number field k and let $\xi \in \text{III}(E/k)[3]$. Then ξ is visible in the jacobian of a genus 2 curve C/k .*

Proof. Let $\delta \in S^{(3)}(E/k)$ be a cocycle representing ξ . By Proposition 2.4, there is a 3-covering $C_\delta \rightarrow E$ corresponding to δ . According to Remark 2.5, we have that $\text{Ob}(\delta) = 0$ and hence that $C_\delta \subset \mathbb{P}^2$. Let $\Phi \subset \mathbb{P}^2$ be its flex scheme. The construction in Section 4 gives us a pencil of cubics through Φ^* , so we can easily pick a non-singular one with a rational point. It follows from Proposition 4.2 that such a curve is of the form $C_{\lambda(\delta)}$ for some elliptic curve E_2 and some anti-isometry $\lambda : E[3] \rightarrow E_2[3]$.

This places us in the situation of Lemma 3.1, so δ is visible in an abelian surface $A = (E \times E_2)/\Delta$. We have ensured that λ is an anti-isometry, which implies that the surface is principally polarized. As long as we make sure that E, E_2 are non-isogenous (and this is easy given the freedom we have in choosing $C_\lambda(\delta)$) it follows that A is a jacobian. $\square$

Remark 5.2. We could of course state a more general result about visibility of elements $\delta \in H^1(k, E[3])$ with $\text{Ob}(\delta) = 0$ for an elliptic curves E over a perfect field k of characteristic distinct from 2 or 3. Note however that if k is too small, there might not be enough non-isogenous elliptic curves available. The exclusion of fields of characteristic 3 is a serious one, the exclusion of non-perfect fields less so. Most of what we are saying could be generalized to the non-perfect case, basically because for an elliptic curve over any field of characteristic distinct from 3, the multiplication by 3 map is separable. The exclusion of fields of characteristic 2 stems from the fact that the necessary invariant theory in this case is not readily available.

We continue with the construction of the genus 2 curve C . Define the divisor $\Theta := 0_1 \times E_2 + E_1 \times 0_2$ on $E_1 \times E_2$, which gives a principal polarization on $E_1 \times E_2$. Next, consider the set $\mathcal{D}$ of effective divisors on $E_1 \times E_2$ over $\bar{k}$ which are linear equivalent to 3Θ and invariant under Δ . Also consider the set $\mathcal{C}$ of effective divisors C on A over $\bar{k}$ whose pull-back to $E_1 \times E_2$ are linear equivalent to 3Θ and which satisfy $(C \cdot C) = 2$. Frey and Kani show that there exist unique curves $D \in \mathcal{D}$ and $C \in \mathcal{C}$ defined over k which are invariant under multiplication by -1 . Furthermore, because E_1 and E_2 are not isogenous, D and C are irreducible smooth curves of genus 10 and 2 respectively and the natural map $D \rightarrow C$ is unramified of degree 9.

If k is a perfect field of characteristic distinct from 2 or 3, the curves D and C can be explicitly constructed as follows. Embed E_1 in $\mathbb{P}^2$, given by, say $F(x, y, z) = 0$, for a ternary cubic F/k (such an F is readily obtained if E_1 is given by a Weierstrass model). Express E_2 as $G := sP(F) + tQ(F) = 0$ for some $s, t \in k$. This way, we obtain an embedding of $E_1 \times E_2$ in $\mathbb{P}^2 \times \mathbb{P}^2$ given by

$$F(x, y, z) = G(u, v, w) = 0.$$

Moreover, by appealing to Proposition 4.2 we obtain that the curve on this surface given by $xu + yv + zw = 0$ must be the curve D . The genus 2 curve C is the image

of D in $(E_1 \times E_2)/\Delta$.

$$\begin{array}{ccc}
 E_1 \times E_2 & & \\
 \downarrow [3] \times [3] & \searrow & \\
 & (E_1 \times E_2)/\Delta & \\
 & \swarrow & \\
 E_1 \times E_2 & &
 \end{array}$$

The map $[3] \times [3]$ is much more accessible, though. Also observe that the hypersurface $\{xu + yv + zw = 0\} \subset \mathbb{P}^2 \times \mathbb{P}^2$ is only invariant under $\Delta \subset E_1[3] \times E_2[3]$. A little extra work shows that the subgroup of $E_1[3] \times E_2[3]$ under which D is invariant is equal to Δ . Hence, we can find a model of C as a curve on $E_1 \times E_2$ by computing $([3] \times [3])(D)$. This can easily be done via interpolation, as explained below by means of an example.

6. EXAMPLES

Following the first example in [2, Table 1], consider the elliptic curve 681b1 (in Cremona's notation), given by the minimal Weierstrass equation

$$E_1 : y^2 + xy = x^3 + x^2 - 1154x - 15345.$$

It turns out that the plane cubic curve

$$C_1 : x^3 + 5x^2y + 5x^2z + 2xy^2 + xyz + xz^2 + y^3 - 5y^2z + 2yz^2 + 6z^3 = 0$$

defines an element ξ (up to inverse) of order three in $\text{III}(E_1/\mathbb{Q})$. The contravariants of the cubic above defining C_1 , denoted P_0, Q_0 , are given by

$$\begin{aligned}
 P_0 &= -478x^3 + 2525x^2y + 916x^2z - 1127xy^2 + 29xyz \\
 &\quad - 160xz^2 + 753y^3 - 1228y^2z + 260yz^2 + 301z^3, \\
 Q_0 &= -122314x^3 + 618551x^2y + 191092x^2z - 271157xy^2 - 7825xyz \\
 &\quad - 28120xz^2 + 184011y^3 - 264916y^2z + 55892yz^2 + 73663z^3.
 \end{aligned}$$

Now the curve

$$C_2 : 55033P_0 - 235Q_0 = 0$$

has a rational point $[x : y : z] = [10 : 8 : 7]$ and its jacobian is the elliptic curve 681c1, given by the minimal Weierstrass equation

$$E_2 : y^2 + y = x^3 - x^2 + 2.$$

To construct the corresponding genus two curve C such that ξ becomes visible in its jacobian we could now take the curve in $C_1 \times C_2 \subset \mathbb{P}^2 \times \mathbb{P}^2$ with coordinates $([x : y : z], [u : v : w])$ given by the equation $xu + yv + zw = 0$, and take its image under $C_1 \times C_2 \rightarrow E_1 \times E_2$, since this is a twist of $[3] \times [3] : E_1 \times E_2 \rightarrow E_1 \times E_2$ anyway. We will follow Section 5 more closely. Obviously, E_1 is given by $F = 0$ if we define

$$F := y^2z + xyz - (x^3 + x^2z - 1154xz^2 - 15345z^3).$$

The contravariants of the ternary cubic F are given by

$$\begin{aligned} P &= -2308x^3 + 3462x^2y - 5x^2z - 275056xy^2 + 5xyz \\ &\quad + 6xz^2 + 136951y^3 + 13853y^2z - 3yz^2, \\ Q &= -725020x^3 + 1087530x^2y + 27721x^2z - 65861608xy^2 - 27721xyz \\ &\quad - 30xz^2 + 32749549y^3 + 3217559y^2z + 15yz^2 + 24z^3. \end{aligned}$$

Write $j(s, t)$ for the j -invariant of the curve given by $sP + tQ = 0$. The j -invariant of E_2 equals $-4096/2043$ and the equation $j(s, t) = -4096/2043$ has exactly one solution in $\mathbb{P}^1(\mathbb{Q})$, namely $[s : t] = [55033 : -235]$ (compare with the definition of C_2). This gives us a new model for E_2 , namely

$$E_2 : 55033P - 235Q = 0.$$

We consider the surface $E_1 \times E_2$ embedded in $\mathbb{P}^2 \times \mathbb{P}^2$ as

$$F(x, y, z) = 0, \quad 55033P(u, v, w) - 235Q(u, v, w) = 0.$$

Now D is simply the curve on this surface given by

$$xu + yv + zw = 0.$$

The image of D under multiplication by 3 on $E_1 \times E_2$ is the genus two curve C . Using the defining properties of C from Section 5 (such as the invariance under multiplication by -1), we get that as a curve on $E_1 \times E_2$ it must be of the form

$$axu + byv + czw + dxw + ezu = 0$$

for some $a, b, c, d, e \in \mathbb{Q}$. Now we simply generate 4 points on C (over a number field), compute the image under multiplication by 3 of these points and solve for a, b, c, d, e . If the dimension of the solution space is greater than 1, we must of course add points (or take 4 better ones) so that the solution space becomes 1-dimensional. This gives us our equation for C . By a linear change of the u, v, w coordinates we can change the model for E_2 back to the original minimal Weierstrass model. Thus, the model for $E_1 \times E_2$ embedded in $\mathbb{P}^2 \times \mathbb{P}^2$ is

$$\begin{aligned} E_1 : y^2z + xyz &= x^3 + x^2z - 1154xz^2 - 15345z^3, \\ E_2 : v^2w + vw^2 &= u^3 - u^2w + 2w^3 \end{aligned}$$

and C is the curve on this surface given by

$$4xu - 155zu + xv + 2yv - 40xw + yw + 1314zw = 0.$$

Hyperelliptic models for C are

$$\begin{aligned} Y^2 + (X + 1)Y &= 3X^5 + 5X^4 + X^3 - 8X^2 - 5X + 2 \text{ or} \\ Y^2 &= (3X - 1)(X + 1)(4X^3 + 4X^2 - 9). \end{aligned}$$

Next, consider the elliptic curve 2006e1, given by the minimal Weierstrass equation

$$E_1 : y^2 + xy = x^3 + x^2 - 58293654x - 171333232940.$$

It turns out that the plane cubic curve

$$C_1 : 20x^3 + 44x^2y + 21x^2z - 77xy^2 + 71xyz + 44xz^2 + 31y^3 + 3y^2z + 150yz^2 + z^3 = 0$$

defines an element ξ (up to inverse) of order three in $\text{III}(E_1/\mathbb{Q})$. In the sixth example in [2, Table 1] the elliptic curve E_2 which ‘explains’ $\text{III}(E_1/\mathbb{Q})$ is 2006d1.

However, for this choice of E_2 , there only exists an isometry between $E_1[3]$ and $E_2[3]$ and not an anti-isometry. The corresponding abelian surface $(E_1 \times E_2)/\Delta$ visualizing ξ will not be the jacobian of a genus 2 curve. If instead we take for E_2 the elliptic curve 6018c1, then we do have an anti-isometry between $E_1[3]$ and $E_2[3]$. Following the same route as in the first example, we find that ξ is visible in the jacobian of the genus 2 curve C with hyperelliptic models

$$\begin{aligned} Y^2 + (X^2 + X)Y &= -9675X^6 - 94041X^5 - 914X^4 + 1301674X^3 - 352310X^2 \\ &\quad - 2071181X - 945269 \text{ or} \\ Y^2 &= 43(2X + 13)(18X^2 - 81X + 89)(25X^3 + 193X^2 + 224X + 76). \end{aligned}$$

REFERENCES

- [1] J. E. Cremona, T. A. Fisher, C. O’Neil, D. Simon, and M. Stoll, *Explicit n -descent on elliptic curves. I. Algebra*, J. Reine Angew. Math. **615** (2008), 121–155.
- [2] John E. Cremona and Barry Mazur, *Visualizing elements in the Shafarevich-Tate group*, Experiment. Math. **9** (2000), no. 1, 13–28.
- [3] B. Mazur, *Visualizing elements of order three in the Shafarevich-Tate group*, Asian J. Math. **3** (1999), no. 1, 221–232. Sir Michael Atiyah: a great mathematician of the twentieth century.
- [4] Catherine O’Neil, *The period-index obstruction for elliptic curves*, J. Number Theory **95** (2002), no. 2, 329–339.
- [5] Tom Fisher, *The Hessian of a genus one curve*, arXiv: math/0610403 (2006), available at <http://lanl.arxiv.org/abs/math/0610403>.
- [6] George Salmon, *A treatise on the higher plane curves*, Third edition, Hodges, Foster, and Figgis, Grafton Street, Dublin, 1879.
- [7] Gerhard Frey and Ernst Kani, *Curves of genus 2 covering elliptic curves and an arithmetical application*, Arithmetic algebraic geometry (Texel, 1989), 1991, pp. 153–176.

DEPARTMENT OF MATHEMATICS, SIMON FRASER UNIVERSITY, BURNABY, BC, CANADA
E-mail address: `nbruin@sfu.ca`
E-mail address: `sdahmen@irmacs.sfu.ca`